## IMMUNIZATION SCHEDULE – CENTRE PEDIATRIC ASSOCIATES MASSACHUSETTS AND AAP REQUIREMENTS – 2025

| IMMUNIZATIONS      |                                 |         |                   |          |                   |                  |                                    |                        |                       |
|--------------------|---------------------------------|---------|-------------------|----------|-------------------|------------------|------------------------------------|------------------------|-----------------------|
|                    | DTAP/HIB<br>POLIO/HEP B<br>TDAP | HEP B 1 | MMR/VZV           | PCV20    | НЕР А             | FLU <sup>2</sup> | MENINGITIS <sup>3</sup>            | ROTAVIRUS <sup>4</sup> | HPV                   |
| 1 MO               |                                 | Нер В   |                   |          |                   |                  |                                    |                        |                       |
| 2 MO <sup>5</sup>  | Vaxelis<br>(DTaP/HIB/IPV/HBV)   |         |                   | PCV20 #1 |                   |                  |                                    | Rotateq #1             |                       |
| 4 MO               | Vaxelis<br>(DTaP/HIB/IPV/HBV)   |         |                   | PCV20 #2 |                   |                  |                                    | Rotateq #2             |                       |
| 6 MO               | Vaxelis<br>(DTaP/HIB/IPV/HBV)   |         | Note <sup>6</sup> | PCV20 #3 | Note <sup>7</sup> | FLU              |                                    | Rotateq #3             |                       |
| 12 MO              |                                 |         | MMR &<br>VZV      |          | Нер А             | FLU              |                                    |                        |                       |
| 15 MO              | Pentacel (DTaP/HIB/IPV)         |         |                   | PCV20 #4 |                   | FLU              |                                    |                        |                       |
| 18 MO              |                                 |         |                   |          | Нер А             | FLU              |                                    |                        |                       |
| 4 YRS              | Quadracel<br>(DTaP/IPV)         |         | ProQuad<br>(MMRV) |          |                   | FLU              |                                    |                        |                       |
| 11 YRS             | TDAP <sup>8</sup>               |         |                   |          |                   | FLU              | Menquadfi                          |                        | Gardasil <sup>9</sup> |
| 16 YRS             |                                 |         |                   |          |                   | FLU              | Menquadfi<br>Bexsero <sup>10</sup> |                        |                       |
| 18 YRS/<br>College |                                 |         |                   |          |                   | FLU              |                                    |                        |                       |

## MUST BE OF AGE AT THESE APPOINTMENTS TO RECEIVE VACCINES

<sup>1</sup> HEP B: If not administered at birth, give initial dose at newborn visit; administer 2<sup>nd</sup> dose at 1 mo if mom is HEP B positive

<sup>2</sup> FLU: Give >6 mo September to June; if under 8 yo AND if there is only 1 previously documented dose, then will need two for current season

<sup>3</sup> MENINGITIS: 11 yo dose, then booster at 16 yo; if 1<sup>st</sup> dose at 13-15 yo, then booster at 16-18 yo; if 1<sup>st</sup> dose age 16, no

booster needed – see CDC recommended child/adolescent immunization schedule age 0-18 yo regarding pt with high-risk condition or immunocompromise

<sup>4</sup> ROTATEQ: Maximum age to start series is 15 wo, maximum age to complete series is 8 mo

<sup>5</sup> 2 MO: Earliest age for 2 mo vaccines is 6 wo; if given at 6 wo, you may give next series of vaccines 4 weeks later at 10 wo

<sup>6</sup> MMR: Can receive at 6 mo for travel purposes, will still need 2-dose series after 12 mo

<sup>7</sup> HEP A: ≥12 mo: 2 doses 6 mo apart, can be given at 6 mo for travel, will still need 2-dose series after 12 mo

<sup>8</sup> TDAP: ≥12 mo: 2 doses 6 mo apart, can be given at 6 mo for travel, will still need 2-dose series after 12 mo

11 yo visit, booster every 10 years, administer for puncture wound with metal if >5 years since last dose

<sup>9</sup> GARDASIL: If first dose <15 yo, then give 2 doses 6 mo apart; if first dose >15 yo, then give 3 doses at 0, 2, and 6 mo – Gardasil may be started as early as 9 yo

<sup>10</sup> BEXSERO: 2<sup>nd</sup> dose at least 6 mo apart