IMMUNIZATION SCHEDULE – CENTRE PEDIATRIC ASSOCIATES
MASSACHUSETTS AND AAP REQUIREMENTS – 2026

AGE	VAXELIS PENTACEL QUADRACEL TDAP	HIB	MMR/VZV	PCV20	HEP B [1]	HEP A	FLU[2]	MENINGITIS[3]	ROTATEQ[4]	HPV
1 MO					Hep B					
2 MO[5]	Vaxelis (DTaP/HIB/IPV/HBV)			PCV20 #1					Rotateq #1	
4 MO	Vaxelis (DTaP/HIB/IPV/HBV)			PCV20 #2					Rotateq #2	
6 MO	Vaxelis (DTaP/HIB/IPV/HBV)		Note [6]	PCV20 #3		Note [7]	Flu		Rotateq #3	
12 MO			MMR & VZV			Hep A	Flu			
15 MO	Pentacel (DTaP/HIB/IPV)			PCV20 #4			Flu			
18 MO						Hep A	Flu			
4 YRS	Quadracel (DTaP/IPV)		ProQuad (MMRV)				Flu			
11 YRS	TDAP [8]						Flu	Menquadfi		Gardasil [9]
16 YRS							Flu	Menquadfi Bexsero [10]		
18 YRS/ College							Flu			

MUST BE OF AGE AT THESE APPOINTMENTS TO RECEIVE VACCINES

[1] HEP B: If not administered at birth, give initial dose at newborn visit. See CDC guidelines if mom is hep B positive. Dose 2 (Vaxelis #1) must be 4 weeks apart from Dose 1 ***and*** pt must be at least 6 weeks old. Dose 4 must be at least 8 weeks apart from dose 2 and at least 16 weeks after dose 1: minimum age for the final dose is 24 weeks.

[2] FLU: Give >6 mo September to June; if under 8 yo AND if there is only 1 previously documented dose, then will need two for current season. Flumist for patients >2 years old, no asthma/wheezing history, not immunocomprised.

[3] MENINGITIS: 11 yo dose, then booster at 16 yo; if 1st dose at 13-15 yo, then booster at 16-18 yo; if 1st dose age 16, no booster needed – see CDC recommended child/adolescent immunization schedule age 0-18 yo regarding pt with high-risk condition or immunocompromise

[4] ROTATEQ: Maximum age to start series is 15 weeks, maximum age to complete series is 8mo.

[5] 2 MO: Earliest age for 2 mo vaccines is 6 wo; if given at 6 wo, you may give next series of vaccines 4 weeks later at 10 wo

[6] MMR: Can receive at 6 mo for travel purposes, will still need 2-dose series after 12 mo. Must be at least 4 weeks between doses

[7] HEP A: ≥12 mo: 2 doses 6 mo apart; can be given at 6 mo for travel, will still need 2-dose series after 12 mo

[8] TDAP: 11 yo visit, booster every 10 years (booster can be given as early as 8 years), administer for puncture wound with metal if >5 years since last dose.

[9] GARDASIL: If first dose <15 yo, then give 2 doses 6 mo apart; if first dose >15 yo, then give 3 doses at 0, 2, and 6 mo – Gardasil may be started as early as 9 yo

[10] BEXSERO: 2nd dose at least 6 mo apart